

To appear in Adv. in Appl. Math.

A CONNECTION BETWEEN COVERS OF THE INTEGERS AND UNIT FRACTIONS

ZHI-WEI SUN

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China

Received 8 July 2005; accepted 2 February 2006

ABSTRACT. For integers a and $n > 0$, let $a(n)$ denote the residue class $\{x \in \mathbb{Z} : x \equiv a \pmod{n}\}$. Let A be a collection $\{a_s(n_s)\}_{s=1}^k$ of finitely many residue classes such that A covers all the integers at least m times but $\{a_s(n_s)\}_{s=1}^{k-1}$ does not. We show that if n_k is a period of the covering function $w_A(x) = |\{1 \leq s \leq k : x \in a_s(n_s)\}|$ then for any $r = 0, \dots, n_k - 1$ there are at least m integers in the form $\sum_{s \in I} 1/n_s - r/n_k$ with $I \subseteq \{1, \dots, k-1\}$.

MSC: primary 11B25; secondary 11B75, 11D68.

1. INTRODUCTION

For an integer a and a positive integer n , we use $a(n)$ to denote the residue class $\{x \in \mathbb{Z} : x \equiv a \pmod{n}\}$. For a finite system

$$A = \{a_s(n_s)\}_{s=1}^k \tag{1}$$

of residue classes, the function $w_A : \mathbb{Z} \rightarrow \{0, 1, \dots\}$ given by

$$w_A(x) = |\{1 \leq s \leq k : x \in a_s(n_s)\}|$$

is called the *covering function* of A . Clearly $w_A(x)$ is periodic modulo the least common multiple N_A of the moduli $n_1, \dots, n_k$, and it is easy to verify the following well-known equality:

$$\frac{1}{N_A} \sum_{x=0}^{N_A-1} w_A(x) = \sum_{s=1}^k \frac{1}{n_s}.$$

E-mail address: zwsun@nju.edu.cn *URL:* <http://pweb.nju.edu.cn/zwsun>

Supported by the National Science Fund for Distinguished Young Scholars (no. 10425103) and a Key Program of NSF (no. 10331020) in China.

As in [7] we call $m(A) = \min_{x \in \mathbb{Z}} w_A(x)$ the *covering multiplicity* of A . For example,

$$B = \{0(2), 0(3), 1(4), 5(6), 7(12)\}$$

has covering multiplicity $m(B) = 1$, because the covering function is periodic modulo $N_B = 12$, and

$$w_B(x) = \begin{cases} 1 & \text{if } x \in \{1, 2, 3, 4, 7, 8, 10, 11\}, \\ 2 & \text{if } x \in \{0, 5, 6, 9\}. \end{cases}$$

Let m be a positive integer. If $w_A(x) \geq m$ for all $x \in \mathbb{Z}$, then we call A an m -cover of the integers, and in this case we have the well-known inequality $\sum_{s=1}^k 1/n_s \geq m$. (The term “1-cover” is usually replaced by the word “cover”.) If A is an m -cover of the integers but $A_t = \{a_s(n_s)\}_{s \in [1, k] \setminus \{t\}}$ is not (where $[a, b] = \{x \in \mathbb{Z} : a \leq x \leq b\}$ for $a, b \in \mathbb{Z}$), then we say that A forms an m -cover of the integers with $a_t(n_t)$ *irredundant*. (For example, $\{0(2), 1(2), 2(3)\}$ is a cover of the integers in which $2(3)$ is redundant while $0(2)$ and $1(2)$ are irredundant.) If $w_A(x) = m$ for all $x \in \mathbb{Z}$, then A is said to be an *exact m -cover* of the integers, and in this case we have the equality $\sum_{s=1}^k 1/n_s = m$. By a graph-theoretic argument, M. Z. Zhang [12] showed that if $m > 1$ then there are infinitely many exact m -covers of the integers each of which cannot be split into an n -cover and an $(m - n)$ -cover of the integers with $0 < n < m$.

Covers of the integers by residue classes were first introduced by P. Erdős (cf. [1]) in the 1930s, who observed that the system B mentioned above is a cover of the integers with distinct moduli. The topic of covers of the integers has been an active one in combinatorial number theory (cf. [3, 4]), and many surprising applications have been found (see, e.g., [1, 2, 9, 11]). The so-called m -covers and exact m -covers of the integers were systematically studied by the author in the 1990s.

Concerning the cover B given above one can easily check that

$$\begin{aligned} & \left\{ \sum_{n \in S} \frac{1}{n} : S \subseteq \{2, 3, 4, 6, 12\} \right\} \\ &= \left\{ 0, \frac{1}{12}, \dots, \frac{11}{12} \right\} \cup \left\{ 1 + \frac{r}{12} : r = 0, 1, 2, 3, 4 \right\}. \end{aligned}$$

This suggests that for a general m -cover (1) of the integers we should investigate the set $\{\sum_{s \in I} 1/n_s : I \subseteq [1, k]\}$.

In this paper we establish the following new connection between covers of the integers and unit fractions.

Theorem 1. *Let $A = \{a_s(n_s)\}_{s=1}^k$ be an m -cover of the integers with the residue class $a_k(n_k)$ irredundant. If the covering function $w_A(x)$ is*

periodic modulo n_k , then for any $r = 0, \dots, n_k - 1$ we have

$$\left| \left\{ \left\lfloor \sum_{s \in I} \frac{1}{n_s} \right\rfloor : I \subseteq [1, k-1] \text{ and } \left\{ \sum_{s \in I} \frac{1}{n_s} \right\} = \frac{r}{n_k} \right\} \right| \geq m, \quad (2)$$

where $\lfloor \alpha \rfloor$ and $\{\alpha\}$ denote the integral part and the fractional part of a real number α respectively.

Note that n_k in Theorem 1 needn't be the largest modulus among $n_1, \dots, n_k$. In the case $m = 1$ and $n_k = N_A$, Theorem 1 is an easy consequence of [5, Theorem 1] as observed by the author's twin brother Z. H. Sun. When $w_A(x) = m$ for all $x \in \mathbb{Z}$, the author [6] even proved the following stronger result:

$$\left| \left\{ I \subseteq [1, k-1] : \sum_{s \in I} \frac{1}{n_s} = \frac{a}{n_k} \right\} \right| \geq \binom{m-1}{\lfloor a/n_k \rfloor} \quad \text{for all } a = 0, 1, \dots$$

Given an m -cover $\{a_s(n_s)\}_{s=1}^k$ of the integers with $a_k(n_k)$ irredundant, by refining a result in [7] the author can show that there exists a real number $0 \leq \alpha < 1$ such that (2) with r/n_k replaced by $(\alpha + r)/n_k$ holds for every $r = 0, \dots, n_k - 1$.

Here we mention two local-global results related to Theorem 1.

(a) (Z. W. Sun [5]) $\{a_s(n_s)\}_{s=1}^k$ forms an m -cover of the integers if it covers $|\{\sum_{s \in I} 1/n_s\} : I \subseteq [1, k]\}|$ consecutive integers at least m times.

(b) (Z. W. Sun [10]) $\{a_s(n_s)\}_{s=1}^k$ is an exact m -cover of the integers if it covers $|\bigcup_{s=1}^k \{r/n_s : r \in [0, n_s - 1]\}|$ consecutive integers exactly m times.

Corollary 1. *Suppose that the covering function of $A = \{a_s(n_s)\}_{s=1}^k$ has a positive integer period n_0 . If there is a unique $a_0 \in [0, n_0 - 1]$ such that $w_A(a_0) = m(A)$, then for any $D \subseteq \mathbb{Z}$ with $|D| = m(A)$ we have*

$$\left\{ \left\{ \sum_{s \in I} \frac{1}{n_s} \right\} : I \subseteq [1, k] \text{ and } \left\lfloor \sum_{s \in I} \frac{1}{n_s} \right\rfloor \notin D \right\} \supseteq \left\{ \frac{r}{n_0} : r \in [0, n_0 - 1] \right\}.$$

Proof. Let $m = m(A) + 1$. Clearly $A' = \{a_s(n_s)\}_{s=0}^k$ forms an m -cover of the integers with $a_0(n_0)$ irredundant. As $w_{A'}(x) - w_A(x)$ is the characteristic function of $a_0(n_0)$, $w_{A'}(x)$ is also periodic mod n_0 . Applying Theorem 1 we immediately get the desired result. $\square$

We will prove Theorem 1 in Section 3 with help from some lemmas given in the next section.

2. SEVERAL LEMMAS

Lemma 1. *Let (1) be a finite system of residue classes with $m(A) = m$, and let $m_1, \dots, m_k$ be any integers. If $f(x_1, \dots, x_k)$ is a polynomial with coefficients in the complex field $\mathbb{C}$ and $\deg f \leq m$, then for any $z \in \mathbb{Z}$ we have*

$$\begin{aligned} & \sum_{I \subseteq [1, k]} (-1)^{|I|} f(\llbracket 1 \in I \rrbracket, \dots, \llbracket k \in I \rrbracket) e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s} \\ &= (-1)^k c(I_z) \prod_{s \in [1, k] \setminus I_z} \left(e^{2\pi i (a_s - z) m_s / n_s} - 1 \right), \end{aligned} \quad (3)$$

where $\llbracket s \in I \rrbracket$ takes 1 or 0 according as $s \in I$ or not, $I_z = \{1 \leq s \leq k : z \in a_s(n_s)\}$, and $c(I_z) = [\prod_{s \in I_z} x_s] f(x_1, \dots, x_k)$ is the coefficient of the monomial $\prod_{s \in I_z} x_s$ in $f(x_1, \dots, x_k)$.

Proof. Write $f(x_1, \dots, x_k) = \sum_{j_1, \dots, j_k \geq 0} c_{j_1, \dots, j_k} x_1^{j_1} \cdots x_k^{j_k}$. Observe that

$$\begin{aligned} & \sum_{I \subseteq [1, k]} (-1)^{|I|} f(\llbracket 1 \in I \rrbracket, \dots, \llbracket k \in I \rrbracket) e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s} \\ &= \sum_{\substack{j_1, \dots, j_k \geq 0 \\ j_1 + \dots + j_k \leq m}} c_{j_1, \dots, j_k} \sum_{I \subseteq [1, k]} \left(\prod_{s=1}^k \llbracket s \in I \rrbracket^{j_s} \times (-1)^{|I|} e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s} \right) \\ &= \sum_{\substack{j_1, \dots, j_k \geq 0 \\ j_1 + \dots + j_k \leq m}} c_{j_1, \dots, j_k} \sum_{J(j_1, \dots, j_k) \subseteq I \subseteq [1, k]} (-1)^{|I|} e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s}, \end{aligned}$$

where $J(j_1, \dots, j_k) = \{1 \leq s \leq k : j_s \neq 0\}$.

Let z be any integer. If $I_z \not\subseteq J(j_1, \dots, j_k)$, then

$$\sum_{J(j_1, \dots, j_k) \subseteq I \subseteq [1, k]} (-1)^{|I|} e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s} = 0$$

since

$$\begin{aligned} & \sum_{I \subseteq [1, k] \setminus J(j_1, \dots, j_k)} (-1)^{|I|} e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s} \\ &= \prod_{s \in [1, k] \setminus J(j_1, \dots, j_k)} \left(1 - e^{2\pi i (a_s - z) m_s / n_s} \right) = 0. \end{aligned}$$

If $j_1, \dots, j_k$ are nonnegative integers with $j_1 + \dots + j_k \leq m$ and $I_z \subseteq J(j_1, \dots, j_k)$, then

$$m = m(A) \leq |I_z| \leq |J(j_1, \dots, j_k)| \leq j_1 + \dots + j_k \leq m;$$

hence $I_z = J(j_1, \dots, j_k)$ and $j_s = 1$ for all $s \in I_z$.

Combining the above we find that the left-hand side of (3) coincides with

$$\begin{aligned} & c(I_z) \sum_{I_z \subseteq I \subseteq [1, k]} (-1)^{|I|} e^{2\pi i \sum_{s \in I} (a_s - z) m_s / n_s} \\ &= c(I_z) (-1)^{|I_z|} e^{2\pi i \sum_{s \in I_z} (a_s - z) m_s / n_s} \prod_{s \in [1, k] \setminus I_z} \left(1 - e^{2\pi i (a_s - z) m_s / n_s}\right) \\ &= (-1)^k c(I_z) \prod_{s \in [1, k] \setminus I_z} \left(e^{2\pi i (a_s - z) m_s / n_s} - 1\right). \end{aligned}$$

This proves the desired (3). $\square$

Lemma 2. *Let (1) be an m -cover of the integers with $a_k(n_k)$ irredundant, and let $m_1, \dots, m_{k-1}$ be positive integers. Then, for any $0 \leq \alpha < 1$ we have $C_0(\alpha) = \dots = C_{n_k-1}(\alpha)$, where $C_r(\alpha)$ (with $r \in [0, n_k - 1]$) denotes the sum*

$$\sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} m_s / n_s\} = (\alpha + r) / n_k}} (-1)^{|I|} \binom{\lfloor \sum_{s \in I} m_s / n_s \rfloor}{m-1} e^{2\pi i \sum_{s \in I} (a_s - a_k) m_s / n_s}.$$

Proof. This follows from [7, Lemma 2]. $\square$

Lemma 3. *Let (1) be an m -cover of the integers with $a_k(n_k)$ irredundant. Suppose that n_k is a period of the covering function $w_A(x)$. Then, for any $z \in a_k(n_k)$ we have*

$$\prod_{s \in [1, k] \setminus I_z} \left(1 - e^{2\pi i (a_s - z) / n_s}\right) = \prod_{s \in I_z} n_s \times \prod_{t=1}^{n_k} \left(1 - e^{2\pi i (t - a_k) / n_k}\right)^{w_A(t) - m}$$

where $I_z = \{1 \leq s \leq k : z \in a_s(n_s)\}$.

Proof. Since $a_k(n_k)$ is irredundant, we have $w_A(z_0) = m$ for some $z_0 \in a_k(n_k)$. As the covering function of A is periodic mod n_k , $|I_z| = w_A(z) = m$ for all $z \in a_k(n_k)$.

Now fix $z \in a_k(n_k)$. Since $w_A(x)$ is periodic modulo n_k , by [8, Lemma 2.1] we have the identity

$$\prod_{s=1}^k \left(1 - y^{N/n_s} e^{2\pi i a_s / n_s}\right) = \prod_{t=1}^{n_k} \left(1 - y^{N/n_k} e^{2\pi i t / n_k}\right)^{w_A(t)},$$

where $N = N_A$ is the least common multiple of $n_1, \dots, n_k$. Putting $y = r^{1/N} e^{-2\pi i z/N}$ where $r \geq 0$, we then get that

$$\prod_{s=1}^k \left(1 - r^{1/n_s} e^{2\pi i(a_s - z)/n_s}\right) = \prod_{t=1}^{n_k} \left(1 - r^{1/n_k} e^{2\pi i(t - z)/n_k}\right)^{w_A(t)}.$$

Therefore

$$\begin{aligned} & \prod_{s \in [1, k] \setminus I_z} \left(1 - e^{2\pi i(a_s - z)/n_s}\right) \\ &= \lim_{r \rightarrow 1} \prod_{s \in [1, k] \setminus I_z} \left(1 - r^{1/n_s} e^{2\pi i(a_s - z)/n_s}\right) \\ &= \lim_{r \rightarrow 1} \frac{\prod_{t=1}^{n_k} (1 - r^{1/n_k} e^{2\pi i(t - a_k)/n_k})^{w_A(t)}}{\prod_{s \in I_z} (1 - r^{1/n_s} e^{2\pi i(a_s - z)/n_s})} \\ &= \lim_{r \rightarrow 1} \prod_{s \in I_z} \frac{1 - r}{1 - r^{1/n_s}} \times \lim_{r \rightarrow 1} \frac{\prod_{t=1}^{n_k} (1 - r^{1/n_k} e^{2\pi i(t - a_k)/n_k})^{w_A(t)}}{(1 - r)^m} \\ &= \prod_{s \in I_z} n_s \times \lim_{r \rightarrow 1} \prod_{t=1}^{n_k} (1 - r^{1/n_k} e^{2\pi i(t - a_k)/n_k})^{w_A(t) - m}, \end{aligned}$$

and hence the desired result follows. $\square$

3. PROOF OF THEOREM 1

In the case $k = 1$, we must have $m = 1$ and $n_k = 1$; hence the required result is trivial. Below we assume that $k > 1$.

Let $r_0 \in [0, n_k - 1]$ and $D = \{d_n + r_0/n_k : n \in [1, m - 1]\}$, where $d_1, \dots, d_{m-1}$ are $m - 1$ distinct nonnegative integers. (If $m = 1$ then we set $D = \emptyset$.) We want to show that there exists an $I \subseteq [1, k - 1]$ such that $\{\sum_{s \in I} 1/n_s\} = r_0/n_k$ and $\sum_{s \in I} 1/n_s \notin D$.

Define

$$f(x_1, \dots, x_{k-1}) = \prod_{d \in D} \left(\frac{x_1}{n_1} + \dots + \frac{x_{k-1}}{n_{k-1}} - d \right).$$

(An empty product is regarded as 1.) Then $\deg f = |D| = m - 1$. For any $z \in a_k(n_k)$, the set $I_z = \{1 \leq s \leq k : z \in a_s(n_s)\}$ has cardinality m since $a_k(n_k)$ is irredundant and $w_A(x)$ is periodic mod n_k . Observe that the coefficient

$$c_z = \left[\prod_{s \in I_z \setminus \{k\}} x_s \right] f(x_1, \dots, x_{k-1}) = \left[\prod_{s \in I_z \setminus \{k\}} x_s \right] \left(\sum_{s=1}^{k-1} \frac{x_s}{n_s} \right)^{m-1}$$

coincides with $(m-1)!/\prod_{s \in I_z \setminus \{k\}} n_s$ by the multinomial theorem. For $I \subseteq [1, k-1]$ we set

$$v(I) = f(\llbracket 1 \in I \rrbracket, \dots, \llbracket k-1 \in I \rrbracket).$$

As $|\{1 \leq s \leq k-1 : x \in a_s(n_s)\}| \geq \deg f$ for all $x \in \mathbb{Z}$, in view of Lemmas 1 and 3 we have

$$\begin{aligned} & \sum_{I \subseteq [1, k-1]} (-1)^{|I|} v(I) e^{2\pi i \sum_{s \in I} (a_s - z)/n_s} \\ &= (-1)^{k-1} c_z \prod_{s \in [1, k-1] \setminus I_z} \left(e^{2\pi i (a_s - z)/n_s} - 1 \right) \\ &= \frac{(-1)^{m-1} (m-1)!}{\prod_{s \in I_z \setminus \{k\}} n_s} \prod_{s \in I_z} n_s \times \prod_{t=1}^{n_k} \left(1 - e^{2\pi i (t - a_k)/n_k} \right)^{w_A(t) - m} = C, \end{aligned}$$

where C is a nonzero constant not depending on $z \in a_k(n_k)$.

By the above,

$$\begin{aligned} N_A C &= \sum_{x=0}^{N_A-1} \sum_{I \subseteq [1, k-1]} (-1)^{|I|} v(I) e^{2\pi i \sum_{s \in I} (a_s - a_k - n_k x)/n_s} \\ &= \sum_{I \subseteq [1, k-1]} (-1)^{|I|} v(I) e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s} \sum_{x=0}^{N_A-1} e^{-2\pi i x \sum_{s \in I} n_k/n_s} \end{aligned}$$

and hence

$$C = \sum_{\substack{I \subseteq [1, k-1] \\ n_k \sum_{s \in I} 1/n_s \in \mathbb{Z}}} (-1)^{|I|} v(I) e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s} = \sum_{r=0}^{n_k-1} C_r,$$

where

$$C_r = \sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} 1/n_s\} = r/n_k}} (-1)^{|I|} \prod_{d \in D} \left(\sum_{s \in I} \frac{1}{n_s} - d \right) e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s}.$$

Let $r \in [0, n_k - 1]$. Write

$$P_r(x) = \prod_{d \in D} \left(x + \frac{r}{n_k} - d \right) = \sum_{n=0}^{m-1} c_{n,r} \binom{x}{n}$$

where $c_{n,r} \in \mathbb{C}$. By comparing the leading coefficients, we find that $c_{m-1,r} = (m-1)!$. Observe that

$$\begin{aligned}
C_r &= \sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} 1/n_s\} = r/n_k}} (-1)^{|I|} P_r \left(\left\lfloor \sum_{s \in I} \frac{1}{n_s} \right\rfloor \right) e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s} \\
&= \sum_{n=0}^{m-1} c_{n,r} \sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} 1/n_s\} = r/n_k}} (-1)^{|I|} \binom{\lfloor \sum_{s \in I} 1/n_s \rfloor}{n} e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s} \\
&= c_{m-1,r} \sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} 1/n_s\} = r/n_k}} (-1)^{|I|} \binom{\lfloor \sum_{s \in I} 1/n_s \rfloor}{m-1} e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s};
\end{aligned}$$

in taking the last step we note that if $0 \leq n < m-1$ then

$$\sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} 1/n_s\} = r/n_k}} (-1)^{|I|} \binom{\lfloor \sum_{s \in I} 1/n_s \rfloor}{n} e^{2\pi i \sum_{s \in I} a_s/n_s} = 0$$

by [5, Theorem 1] (since $\{a_s(n_s)\}_{s=1}^{k-1}$ is an $(m-1)$ -cover of the integers). By Lemma 2 and the above,

$$C_r = (m-1)! \sum_{\substack{I \subseteq [1, k-1] \\ \{\sum_{s \in I} 1/n_s\} = 0}} (-1)^{|I|} \binom{\lfloor \sum_{s \in I} 1/n_s \rfloor}{m-1} e^{2\pi i \sum_{s \in I} (a_s - a_k)/n_s}$$

does not depend on $r \in [0, n_k - 1]$.

Combining the above we obtain that

$$n_k C_{r_0} = \sum_{r=0}^{n_k-1} C_r = C \neq 0.$$

So there is an $I \subseteq [1, k-1]$ for which $\{\sum_{s \in I} 1/n_s\} = r_0/n_k$, $\sum_{s \in I} 1/n_s \notin D$ and hence $\lfloor \sum_{s \in I} 1/n_s \rfloor \notin \{d_n : n \in [1, m-1]\}$. This concludes our proof.

REFERENCES

1. P. Erdős, On integers of the form $2^k + p$ and some related problems, *Summa Brasil. Math.* 2(1950) 113–123.
2. M. Filaseta, Coverings of the integers associated with an irreducibility theorem of A. Schinzel, in: M. A. Bennett, B. C. Berndt, N. Boston, H. G. Diamond, A. J. Hildebrand, W. Philipp (Eds.), *Number Theory for the Millennium* (Urbana, IL, 2000), vol. II, pp. 1-24, A K Peters, Natick, MA, 2002.

3. R. K. Guy, *Unsolved Problems in Number Theory*, third ed., Springer, New York, 2004 (Sections F13 and F14).
4. Š. Porubský and J. Schönheim, Covering systems of Paul Erdős: Past, present and future, in: G. Halász, L. Lovász, M. Simonvits, V. T. Sós (Eds.), *Paul Erdős and his Mathematics, I*, Bolyai Soc. Math. Stud., vol. 11, 2002, pp. 581–627.
5. Z. W. Sun, Covering the integers by arithmetic sequences, *Acta Arith.* 72(1995) 109–129.
6. Z. W. Sun, Exact m -covers and the linear form $\sum_{s=1}^k x_s/n_s$, *Acta Arith.* 81(1997) 175–198.
7. Z. W. Sun, On covering multiplicity, *Proc. Amer. Math. Soc.* 127(1999) 1293–1300.
8. Z. W. Sun, On the function $w(x) = |\{1 \leq s \leq k : x \equiv a_s \pmod{n_s}\}|$, *Combinatorica* 23(2003) 681–691.
9. Z. W. Sun, Unification of zero-sum problems, subset sums and covers of $\mathbb{Z}$, *Electron. Res. Announc. Amer. Math. Soc.* 9(2003) 51–60.
10. Z. W. Sun, Arithmetic properties of periodic maps, *Math. Res. Lett.* 11(2004) 187–196.
11. Z. W. Sun, A local-global theorem on periodic maps, *J. Algebra* 293(2005) 506–512.
12. M. Z. Zhang, Irreducible systems of residue classes that cover every integer exactly m times, *Sichuan Daxue Xuebao (Nat. Sci. Ed.)* 28(1991) 403–408.